

PME1.7: RULES FOR SIGNIFICANT FIGURES

In scientific measurements, significant figures are used to indicate the accuracy of a measurement. The last digit in a measurement is often an estimate, a good guess.

All the digits in a measured value, including the last estimated digit, are called significant figures or significant digits.

Consider a series of measurements made with (a) an old wooden ruler without mm marks, (b) a more accurate steel ruler, (c) steel callipers with a Vernier scale.

	Reported Result	Number of Significant Figures in the measurement	Accuracy of measurement	Implied range of possible values
(a)	7 cm	1	To the nearest 1 cm, ie 7 ± 0.5 cm	6.5 cm – 7.5 cm
(b)	7.2 cm	2	To the nearest 0.1 cm, ie 7.2 ± 0.05 cm	7.15 – 7.25 cm
(c)	7.23	3	To the nearest 0.01 cm, ie 7.23 ± 0.005 cm	7.225 – 7.235 cm

The number of Significant Figures in a measured value

Rule 1 Any non-zero digit is significant. The position of a decimal point makes no difference.

<i>Example 1</i>	15.7	3	sig. figs
	157	3	sig. figs
	1.57	3	sig. figs
	2.7942	5	sig. figs

Rule 2 Zeros between numbers **are** significant.

<i>Example 2</i>	1.05	3	sig. figs
	10.51	4	sig. figs
	200.708	6	sig. figs

Rule 3 Zeros at the right hand end of whole numbers are **not** significant, unless otherwise stated.

<i>Example 3</i>	70	1	sig. figs
	2860	3	sig. figs
	15090	4	sig. figs

Rule 4 Zeros at the left hand end of decimal numbers are **not** significant.

<i>Example 4</i>	0.28	2	sig. figs
	0.0039	2	sig. figs
	0.0604	3	sig. figs

Rule 5 Zeros at the right hand end of decimal numbers **are** significant.

<i>Example 5</i>	12.0	3	sig. figs
	0.760	3	sig. figs
	0.48300	5	sig. figs
	2.07090	6	sig. figs

Significant Figures and Scientific Notation


The problems of deciding how many significant figures a value has is simplified by writing its value in Scientific Notation.

Examine the number of digits in the first number below, not in the power of ten.

- | | | | |
|-----|-----------------------------------|------------|--------|
| (a) | $0.0003 = 3 \times 10^{-4}$ | 1 sig fig | Rule 4 |
| (b) | $720000 = 7.2 \times 10^5$ | 2 sig figs | Rule 3 |
| (c) | $660 = 6.6 \times 10^2$ | 2 sig figs | Rule 3 |
| (d) | $660.0 = 6.600 \times 10^2$ | 4 sig figs | Rule 5 |
| (e) | $0.66000 = 6.6000 \times 10^{-1}$ | 5 sig figs | Rule 5 |
| (f) | $808.01 = 8.0801 \times 10^2$ | 5 sig figs | Rule 2 |

Recording measurements and Significant Figures

Consider the scale below.



The measurement x is:

- Certainly greater than 6 and less than 7
- Certainly greater than 6.5 and less than 6.6
- Very probably greater than 6.53 and less than 6.55

The result would be recorded as 6.54 - a value with 3 significant digits. The first two digits are certain and the last is a good estimate.

Writing "6.54" implies " 6.54 ± 0.005 ", ie between 6.535 and 6.545, unless otherwise stated.

Exercise

Write each of the following measurements in Scientific Notation and state the number of significant digits in the value.

- | | | | |
|-------------|------------|-------------|--------------|
| (a) 345 | (b) 17642 | (c) 0.0033 | (d) 0.000306 |
| (e) 870 | (f) 20000 | (g) 140.600 | (h) 710.0 |
| (i) 0.04080 | (j) 0.0050 | | |

Answers

Exercise

- | | | |
|--------------|------------------------|------------|
| (a) 345 | 3.45×10^2 | 3 sig figs |
| (b) 17642 | 1.7642×10^4 | 5 sig figs |
| (c) 0.0033 | 3.3×10^{-3} | 2 sig figs |
| (d) 0.000306 | 3.06×10^{-4} | 3 sig figs |
| (e) 870 | 8.7×10^2 | 2 sig figs |
| (f) 20000 | 2×10^4 | 1 sig fig |
| (g) 140.600 | 1.40600×10^2 | 6 sig figs |
| (h) 710.0 | 7.100×10^2 | 4 sig figs |
| (i) 0.04080 | 4.080×10^{-2} | 4 sig figs |
| (j) 0.0050 | 5.0×10^{-3} | 2 sig figs |